

Table XXVII-3

AGE-ADJUSTED SEER CANCER INCIDENCE RATES^a, 1975-2000

By Primary Cancer Site

All Races, Males and Females

Year of Diagnosis

<u>Age/Site</u>	1975	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
-----------------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Ages 0-14

All Sites

All Races	11.5	12.9	12.5	12.9	12.9	13.9	14.5	14.3	14.2	13.5	15.0	14.2	15.1	13.4	14.9	13.9	13.9	14.6	14.0	15.3	14.1	15.1	
Whites	12.0	13.1	12.5	13.1	13.7	14.4	15.6	14.9	14.6	14.0	15.5	14.2	16.1	13.9	15.1	14.5	14.6	14.9	14.8	16.2	15.1	16.3	
Blacks	8.6	11.1	13.2	11.8	10.0	10.3	10.1	10.9	11.6	11.9	13.7	12.3	11.6	12.0	12.3	12.2	11.2	11.1	11.1	9.8	10.1	11.7	
Bone & Joint	0.5	0.7	0.6	0.6	0.7	0.7	0.8	0.7	0.7	0.6	0.6	0.7	0.9	0.5	0.6	0.8	0.8	0.5	0.9	0.6	0.6	0.6	
Brain & Other nervous	2.3	2.8	2.5	2.4	2.2	2.8	3.0	3.5	3.2	3.3	3.2	3.5	3.5	3.2	3.4	3.2	3.3	3.1	2.7	3.1	3.3	3.4	
Hodgkin lymphoma	0.7	0.5	0.6	0.5	0.7	0.7	0.5	0.6	0.6	0.7	0.6	0.6	0.5	0.6	0.5	0.5	0.3	0.6	0.6	0.6	0.4	0.6	
Kidney & Renal pelvis	0.7	0.8	0.9	1.0	1.2	0.7	0.9	0.9	0.8	1.1	0.8	0.9	0.7	1.0	1.2	0.6	1.0	1.0	1.1	0.7	1.0		
Leukemia	3.3	4.0	4.1	3.8	3.7	4.5	4.6	4.1	4.4	4.1	4.8	4.4	4.6	4.0	4.2	3.6	4.4	4.6	4.2	4.8	4.6	4.6	
Acute lymphocytic	2.2	3.1	3.3	3.0	3.0	3.5	3.5	3.3	3.4	3.3	4.0	3.4	3.5	3.1	3.1	2.9	3.6	3.5	3.5	3.9	3.5	3.5	
Non-Hodgkin lymphoma	1.0	0.7	0.7	0.9	0.9	0.9	0.7	1.0	1.1	0.7	0.7	0.8	0.9	0.8	0.8	1.0	0.9	0.6	0.9	0.9	0.8	1.0	
Soft tissue	0.9	0.9	0.7	0.9	0.9	0.8	0.9	0.9	0.8	0.6	0.9	0.7	1.1	0.7	0.6	0.9	0.9	0.9	1.1	1.0	1.1	0.9	1.0

Ages 0-19

All Sites

All Races	12.9	14.4	14.0	14.5	14.6	15.5	15.9	15.9	15.3	15.3	16.7	15.7	16.4	16.1	16.2	15.6	15.8	16.2	15.7	16.4	15.4	16.8
Whites	13.3	14.8	14.3	15.1	15.4	16.1	17.1	16.7	16.2	15.8	17.4	15.9	17.5	16.9	16.6	16.2	16.7	16.6	16.8	17.4	16.4	18.0
Blacks	10.8	11.0	13.0	11.8	11.2	12.2	10.6	12.5	11.3	13.0	14.0	13.2	12.4	12.5	13.2	13.1	11.9	12.2	11.0	11.3	10.7	12.9
Bone & Joint	0.7	0.9	1.0	0.8	0.8	0.9	1.0	0.9	1.0	0.8	0.9	0.8	1.0	0.9	0.8	1.0	0.9	0.7	1.2	0.8	0.7	0.8
Brain & Other nervous	2.1	2.5	2.3	2.3	2.2	2.6	2.7	3.2	2.9	3.0	3.0	3.2	3.0	3.2	3.0	2.8	3.0	2.9	2.5	2.7	2.9	3.1
Hodgkin lymphoma	1.5	1.2	1.3	1.5	1.4	1.5	1.4	1.3	1.4	1.3	1.7	1.3	1.4	1.4	1.3	1.3	0.9	1.4	1.3	1.4	1.1	1.1
Kidney & Renal pelvis	0.5	0.6	0.7	0.7	0.9	0.6	0.7	0.7	0.7	0.6	0.9	0.6	0.7	0.6	0.8	0.9	0.5	0.7	0.8	0.8	0.6	0.8
Leukemia	3.0	3.5	3.7	3.4	3.5	4.0	4.0	3.7	3.7	3.6	4.2	3.8	4.1	3.7	3.7	3.4	3.9	3.9	3.7	4.1	4.0	4.1
Acute lymphocytic	1.9	2.5	2.7	2.6	2.6	3.0	2.9	2.8	2.7	2.7	3.4	2.9	3.0	2.6	2.6	2.5	3.0	2.9	2.9	3.3	2.9	3.0
Non-Hodgkin lymphoma	1.0	0.9	1.0	1.0	1.1	1.0	1.1	1.0	1.0	1.0	1.1	1.0	1.1	1.0	0.9	1.2	1.1	1.0	1.1	1.1	0.9	1.3
Soft tissue	0.8	0.9	0.7	1.0	1.0	0.8	0.9	0.9	0.9	0.7	0.9	0.8	1.1	0.7	0.7	0.7	1.1	1.0	1.1	1.0	1.1	1.2

^a SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.